

Adjunctive minocycline for major depressive disorder: Characterisation of peripheral immune-inflammatory profile

Walker AJ¹, Bortolasci C², Mohebbi M^{1,3}, Ashton M^{1,4,5}, Kanchanatawan B⁶, Singh A¹, Ng C⁵, Mahli G^{7,8}, Dowling N⁵, Maes M⁶, Walder K², Berk M^{1,4,9,10,11}, Dean OM^{1,4,9}.



¹IMPACT Strategic Research Centre, School of Medicine, Deakin University, Geelong, VIC, Australia; ²Metabolic Research Unit, School of Medicine, Deakin University, Geelong, VIC, Australia; ³Biostatistics Unit, Faculty of Health, Deakin University; ⁴The Florey Institute for Neuroscience and Mental Health, The University of Melbourne, Parkville, VIC, Australia; ⁵Department of Psychiatry, The Melbourne Clinic, The University of Melbourne, Richmond, VIC, Australia; ⁶Department of Psychiatry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand; ⁷Discipline of Psychiatry, Sydney Medical School, The University of Sydney, Sydney, NSW, Australia; ⁸CADE Clinic, Department of Psychiatry, Royal North Shore Hospital, St. Leonards, NSW, Australia; ⁹Department of Psychiatry, Royal Melbourne Hospital, The University of Melbourne, Parkville, VIC, Australia; ¹⁰Oxygen Youth Health Research Centre, Parkville, VIC, Australia; ¹¹Centre for Youth Mental Health, The University of Melbourne, Parkville, VIC, Australia.

Background

Primary treatments for major depressive disorder (MDD) are useful, however shortfalls remain in both remission and recovery [1].

The identification of biological markers that relate to treatment response may improve understanding of the pathophysiology of MDD and assist with this shortfall [2].

Inflammation is now known to play a role in the pathophysiology of MDD, leading to clinical trials of adjunctive treatments that target these pathways [3].

Minocycline has been shown to modify the immune-inflammatory processes, however its effects on the inflammatory profile of participants with MDD are unknown [4].

Aim: This study aims to address a secondary outcome of a clinical trial by assessing inflammatory biomarkers in collected peripheral blood samples.

Methods

This current study is a secondary analysis of peripheral blood samples collected from patients during a randomised placebo-controlled trial evaluating 12 weeks of adjunctive minocycline (200 mg/day) in addition to treatment as usual (ACTRN12612000283875) [5].

Samples were collected from consenting participants at baseline (week 0), and at the end of treatment (week 12).

Of the 71 participants that completed the trial, at least one serum aliquot was available for 47 of the participants. Placebo group $n=24$ (13f/11m), and the minocycline group $n=23$ (13f/10m).

Samples were assessed for an array of 46 inflammatory analytes including cytokines, chemokines and acute-phase reactants using Myriad RBM's InflammationMAP®

Composite Z-scores were calculated for detectable immune-inflammatory analytes, and statistical analyses were conducted using SPSS Statistical Package version 25.0 by an investigator blinded to treatment.

Preliminary analysis included generation of descriptives, and subsequent Spearman's nonparametric correlation analysis to assess analyte group profile at baseline and endpoint. Measures of depression (MADRS) and anxiety (HAMA) were included to assess treatment response.

Preliminary Results

Spearman's Correlation (ρ) - Baseline

	MADRS	HAMA	AAT	A2M	B2M	BDNF	CRP	C3	CCL11	FVII	FRTN	HAPT	ICAM-1	IL-1 β	IL-1RA	IL-8	IL12p40	IL-15	IL-18	IL-23	MIP-1 β	MMP3	MMP9	MCP1	SCF	RANTES	TIMP1	TNFR2	VCAM-1	VEGF	VDBP	vWF	MINOCYCLINE
MADRS		0.555	0.103	-0.043	0.072	0.317	-0.007	-0.148	-0.083	-0.438	0.082	0.035	0.176	0.219	-0.167	0.214	-0.020	0.049	-0.246	-0.246	-0.028	0.101	0.495	-0.272	-0.290	-0.305	-0.065	0.081	0.208	-0.014	0.176	0.131	
HAMA	0.585		-0.234	0.167	-0.015	0.408	-0.310	0.112	-0.364	-0.562	0.224	-0.302	-0.007	0.140	-0.190	-0.008	-0.405	-0.081	0.060	-0.076	0.047	0.088	0.431	-0.556	-0.441	-0.235	-0.330	-0.036	0.257	0.017	0.063	0.110	
AAT	0.367		-0.011		0.292	0.449	-0.150	0.180	0.236	-0.036	-0.067	-0.359	0.453	0.118	0.209	0.272	0.300	0.077	0.164	-0.417	-0.258	-0.174	0.245	-0.158	0.010	-0.001	0.274	0.447	0.243	0.070	0.190	0.228	0.166
A2M	-0.166	-0.059	0.182	0.468	-0.251	0.102	0.565	-0.518	-0.433	-0.021	0.241	0.268	0.056	-0.096	-0.351	-0.069	-0.252	-0.119	-0.156	-0.367	0.280	0.083	-0.319	-0.354	0.415	0.585	0.212	0.293	-0.101	0.003	0.209		
B2M	0.101	0.116	0.331	0.122	-0.178	0.541	0.207	0.094	-0.054	0.170	0.557	0.537	0.243	0.267	-0.007	0.221	0.593	0.111	0.290	0.084	0.022	0.150	0.110	0.248	0.706	0.678	0.631	-0.119	0.355	0.607			
BDNF	0.085	-0.116	0.315	0.025	0.254		-0.155	-0.005	-0.164	-0.107	-0.227	-0.197	-0.315	0.021	-0.236	0.152	-0.044	0.061	-0.326	0.295	0.039	0.351	0.042	-0.010	0.295	-0.196	0.075	-0.160	0.073	-0.105	0.030		
CRP	-0.168	-0.086	0.007	-0.557	-0.055	0.009		0.314	0.127	0.418	-0.086	0.761	0.406	0.407	0.527	0.531	0.419	0.063	0.334	-0.074	0.097	0.053	0.254	0.156	0.171	0.548	0.373	0.255	-0.034	0.695	0.640		
C3	0.271	-0.160	0.235	-0.286	0.014	0.189	0.545		-0.383	0.082	0.022	0.254	0.160	0.058	0.130	-0.417	0.319	0.044	0.038	0.029	0.111	-0.178	0.138	-0.086	0.097	0.324	0.291	0.146	0.401	0.234	0.427	C3	
CCL11	0.491	0.165	0.328	0.065	0.588	0.450	0.079	0.420		0.396	0.241	0.168	0.000	-0.027	0.131	-0.077	0.322	0.181	0.117	0.241	0.317	0.476	-0.110	0.623	0.421	-0.194	-0.032	0.068	0.083	-0.169	0.241	0.199	
FVII	-0.273	-0.262	-0.121	-0.332	0.022	0.160	0.473	0.636		0.275	0.109	0.331	-0.114	0.160	0.362	0.053	0.600	0.240	0.110	0.278	0.133	0.240	0.151	0.339	0.245	0.240	0.116	0.345	FVII				
FRTN	-0.488	-0.178	-0.062	0.318	0.215	-0.072	0.269	-0.196	-0.355	0.034		0.009	0.415	0.193	-0.071	0.561	-0.115	0.464	0.524	0.511	0.242	0.178	0.298	0.151	0.339	0.245	0.207	0.116	0.345	FRTN			
HAPT	-0.204	-0.192	0.074	-0.491	-0.005	0.003	0.715	0.595	0.168	0.601	0.349	0.183		0.188	0.075	0.275	0.75	0.588	-0.193	0.077	0.384	-0.044	0.256	0.004	0.476	0.179	0.291	0.649	0.416	0.208	-0.011	0.758	0.668
ICAM-1	-0.084	0.131	-0.171	-0.222	0.349	0.270	0.288	-0.054	0.351	0.229	0.188	0.167		0.019	0.001	0.214	0.074	0.153	0.277	0.134	0.302	-0.007	0.148	0.073	0.093	0.022	0.264	0.235	0.355	-0.058	0.194	0.341	
IL-1 β	-0.398	-0.485	0.058	-0.369	0.272	0.580	0.301	0.325	0.320	0.508	0.370	0.362	0.302		0.058	0.527	0.234	0.364	0.646	0.17	0.379	0.070	0.185	0.386	0.359	-0.192	0.273	0.369	0.354	0.198	0.203	0.788	0.716
IL-1RA	0.362	0.244	0.048	-0.324	0.192	0.159	0.597	0.498	0.379	0.390	0.029	0.398	0.442		0.086	-0.077	0.614	0.705	0.272	0.304	0.092	0.111	-0.134	0.238	0.189	-0.063	0.507	0.378	0.074	0.199	0.512	0.504	
IL-8	0.137	0.376	0.153	0.202	-0.054	0.101	0.081	0.178	0.096	0.150	-0.254	-0.048	0.407	-0.291	0.640		0.730	0.554	0.246	0.028	0.267	0.402	-0.020	0.004	0.062	0.513	-0.292	0.178	0.119	0.543	-0.179	0.202	
IL12p40	-0.136	-0.046	-0.503	-0.040	0.027																												